arXiv:1407.1928v1 [physics.ao-ph] 8 Jul 2014

An amplitude equation for long, nonlinear internal waves in a weak shear flow over topography

M. Yu. Trofimov

Il'ichev Pacific Oceanological Institute of the Russian Academy of Sciences, Vladivostok, 690041, Russia

Abstract

A forced, variable coefficients Korteweg-de Vries equation for amplitudes of long, nonlinear internal waves in a stratified shear flow over topography is derived when the magnitude of the basic flow is small. The derivation is done by incorporating the basic flow in the perturbation series of the Taniuti and Wei's reductive perturbation method. Istead of the long-wave Tailor-Goldstein spectral problem, only the usual long-wave spectral problem for rested stratified media and a boundary value problem have to be solved for calculating the coefficients of obtained equation. Explicit expressions for these coefficients are presented.

Known amplitude equations of the Korteweg-de Vries type for long nonlinear internal waves in stratified shear flows are based on the long-wave Tailor-Goldstein spectral problem, which is quadratic in spectral parameter. Contrary to the ordinary self-adjoint spectral problems arising in the absence of basic flow, the quadratic ones are hardly treated numerically because the corresponding algorithms are either complicated or inefficient. In our opinion, this fact restricts the use of such amplitude equations in concrete applications. On the other hand, in many interesting applications the basic flow can be treated as weak. The main idea of our work is to incorporate this flow in the perturbation series of the Taniuti and Wei's reductive perturbation method [1]. The amplitude equation, obtained in this way, turns out to be based on a first-order perturbation of the usual long-wave spectral problem for rested stratified media. This leads to the comparatively simple procedure of calculating the coefficients of this equation.

We shall use the equations of two-dimensional motions of inviscid, incompressible stably stratified fluid, written in the form

$$\begin{aligned}&\rho\left[\partial_t\Delta\Psi + J(\Delta\Psi,\Psi)\right] = \\ &\frac{1}{\beta}\rho_x - \left\{\rho_z\left[\Psi_{tz} + J(\Psi_z,\Psi)\right] + \rho_x\left[\Psi_{tx} + J(\Psi_x,\Psi)\right]\right\}\end{aligned} \tag{1}$$

$$\rho_t + J(\rho,\Psi) = 0 \tag{2}$$

with the boundary conditions

$$\Psi = 0 \quad \text{at} \quad z = 0, \qquad \Psi = 0 \quad \text{at} \quad z = -H(x) \tag{3}$$

where $()_t = \partial_t = \partial/\partial t$, etc., $\Delta = \partial_x^2 + \partial_z^2$, $J(\alpha,\beta) = \alpha_x\beta_z - \alpha_z\beta_x$, Ψ is a stream function, ρ is the density and H describes the bottom topography. The variables are nondimensional, based on a length scale $\bar{h}$ (a typical vertical dimension), a time scale $\bar{N}^{-1}$ (where $\bar{N}$ is a typical value of the Brunt-Väisälä frequency), and a density scale $\bar{\rho}$ (a typical value of the density). The parameter β is $\bar{h}\bar{N}^2 g^{-1}$, where g is the gravity acceleration.

For applying the reductive perturbation method to the equations (1), (2), we introduce a small parameter ϵ and slow variables $\xi = \epsilon\left(\int^x \frac{1}{c}\,dx' - t\right)$, $\eta = \epsilon^3 x$, where $c = c_0 + \epsilon c_1 + \epsilon^2 c_2 + \ldots$ is the phase velocity of the waves to be considered. The amplitude of these waves will be $O(\epsilon^2)$ with the amplitude of basic flow $O(\epsilon)$, so the introduced space-time scales are characteristic for the Korteweg-de Vries theory. The derivatives with respect to x and t are expressed in new variables as follows:

$$\partial_x = \left(\epsilon\frac{1}{c_0} - \epsilon^2\frac{c_1}{c_0^2} + \epsilon^3\frac{c_1^2}{c_0^3} - \epsilon^3\frac{c_2}{c_0^2} + \ldots\right)\partial_\xi + \epsilon^3\partial_\eta\,, \qquad \partial_t = -\epsilon\partial_\xi \tag{4}$$

Expanding ρ and Ψ as $\rho = \rho_0 + \epsilon\rho_1 + \epsilon^2\rho_2 + \ldots$, $\Psi = \epsilon\Psi_1 + \epsilon^2\Psi_2 + \ldots$ and substituting these expansions and Eq. (4) into Eqs. (1), (2) and (3), we obtain a hierarchy of equations at orders of ϵ. We will search a solution satisfying this hierarchy up to the order $O(\epsilon^4)$. In the sequel the bottom topography H is assumed to be dependent only on η.

At $O(\epsilon)$ we have $\rho_{0\xi} = 0$, so ρ_0 does not depend on ξ.

At $O(\epsilon^2)$ we have

$$\begin{aligned}-\rho_0\Psi_{1zz\xi} &= \frac{1}{\beta c_0}\rho_{1\xi} + \rho_{0z}\Psi_{1z\xi} \\ \rho_{1\xi} - \rho_{0z}\frac{1}{c_0}\Psi_{1\xi} &= 0\end{aligned}$$

We suppose that this system is satisfied by the given basic flow which does not depend on ξ. This flow must satisfy the equations (1), (2) up to the order $O(\epsilon^4)$, which gives

$$\rho_{0\eta} = 0, \quad \rho_{1\eta} = 0 \tag{5}$$

$$\rho_{0z}\Psi_{1\eta} = 0 \tag{6}$$

By Eq. (5), ρ_0 and ρ_1 both depend only on z and without loss of generality we take $\rho_1 = 0$. According to Eq. (6), the basic flow has to be pure shear within the pycnocline.

At $O(\epsilon^3)$ we have

$$\begin{aligned} -\rho_0\Psi_{2zz\xi} &= \frac{1}{\beta c_0}\rho_{2\xi} + \rho_{0z}\Psi_{2z\xi} \\ \rho_{2\xi} - \rho_{0z}\frac{1}{c_0}\Psi_{2\xi} &= 0 \end{aligned}$$

As a solution of these equations we take

$$\Psi_2 = c_0(\eta)A(\eta,\xi)Y(\eta,z)\,, \quad \rho_2 = -\rho_{0z}A(\eta,\xi)Y(\eta,z)\,, \tag{7}$$

where A has the meaning of isopycnal elevation amplitude and c_0 together with Y satisfy the usual long-wave spectral problem (see e.g. Grimshaw et al. [2]-[3])

$$c_0^2(\rho_0 Y_z)_z - \frac{1}{\beta}\rho_{0z}Y = 0\,, \qquad Y(0) = Y(-H(\eta)) = 0 \tag{8}$$

with Y normed by

$$-\frac{1}{\beta}\int_{-H(\eta)}^{0}\rho_{0z}Y^2\,dz = 1$$

At $O(\epsilon^4)$, taking into account Eqs. (5) and (6), we have

$$\begin{aligned} &-\rho_0\Psi_{3zz\xi} + \rho_0\frac{1}{c_0}\Psi_{1z}\Psi_{2zz\xi} - \rho_0\frac{1}{c_0}\Psi_{2\xi}\Psi_{1zzz} = \\ &\quad \frac{1}{\beta}\frac{1}{c_0}\rho_{3\xi} - \frac{1}{\beta}\frac{c_1}{c_0^2}\rho_{2\xi} + \rho_{0z}\Psi_{3z\xi} - \frac{1}{c_0}\rho_{0z}\Psi_{1z}\Psi_{2z\xi} + \frac{1}{c_0}\rho_{0z}\Psi_{1zz}\Psi_{2\xi} \\ &\quad \rho_{3\xi} - \frac{1}{c_0}\rho_{2\xi}\Psi_{1z} + \frac{1}{c_0}\rho_{0z}\Psi_{3\xi} - \frac{c_1}{c_0^2}\rho_{0z}\Psi_{2\xi} = 0 \end{aligned}$$

Introducing the horizontal velocity $U = \Psi_{1z}$ and postulating an ansatz for Ψ_3 of the form $c_0(\eta)A(\eta,\xi)W(\eta,z)$, after some calculations we obtain from

these equations a boundary value problem for W

$$\begin{aligned} &c_0^2(\rho_0 W_z)_z - \frac{1}{\beta}\rho_{0z} W = \\ &2c_0 U(\rho_0 Y_z)_z - 2c_1 c_0(\rho_0 Y_z)_z - c_0\rho_0 U_{zz} Y - c_0\rho_{0z} U_z Y \\ &W(0) = W(-H(\eta)) = 0 \end{aligned} \tag{9}$$

It is remarkable that this boundary value problem coincides with the $O(\epsilon)$ part of the long-wave Taylor-Goldstein problem

$$\begin{aligned} &(\epsilon U - c_0 - \epsilon c_1)\left[\rho_0(\epsilon U - c_0 - \epsilon c_1)(Y + \epsilon W)_z\right]_z - \\ &\quad (\epsilon U - c_0 - \epsilon c_1)\left[\rho_0 \epsilon U_z (Y + \epsilon W)\right]_z - \frac{1}{\beta}\rho_{0z}(Y + \epsilon W) = 0 \\ &\\ &Y(0) + \epsilon W(0) = Y(-H(\eta)) + \epsilon W(-H(\eta)) = 0 \end{aligned}$$

which is a perturbation of Eq. (8).

The solvability condition for Eq. (9), which consists in orthogonality of the right-hand side of Eq. (9) to Y in $L^2[-H, 0]$, gives the value for the first order correction to the phase velocity

$$c_1 = c_0^2 \int_{-H}^{0} \rho_0 U (Y_z)^2 \, dz \tag{10}$$

At $O(\epsilon^5)$ we have the system too large to be shown here. It can be reduced to a boundary value problem for $\Psi_{4\xi}$ of the form

$$\begin{aligned} c_0^2(\rho_0 \Psi_{4\xi z})_z - \frac{1}{\beta}\rho_{0z}\Psi_{4\xi} &= \text{R.-H. S.} \\ \Psi_{4\xi}(0) = \Psi_{4\xi}(-H) &= 0 \end{aligned}$$

where R.-H. S. stands for a collection of terms containing A, Y, W, c_0 and c_1. The solvability condition for this problem, which is the same as for Eq. (9), gives an amplitude equation of the Korteweg-de Vries type

$$\alpha A + A_\eta + \beta A A_\xi + \gamma A_\xi + \delta A_{\xi\xi\xi} = \mathcal{F} \tag{11}$$

where

$$\begin{aligned}
\alpha &= \frac{1}{2}\frac{\partial}{\partial\eta}\ln c_0 \\
\beta &= \frac{3}{2}c_0\int_{-H}^{0}\rho_0(Y_z)^3\,dz \\
\gamma &= -2\int_{-H}^{0}\rho_0(U-c_1)W_zY_z\,dz - \int_{-H}^{0}\rho_0U_z(W_zY-WY_z)\,dz + \\
&\quad \frac{1}{c_0}\int_{-H}^{0}\rho_0\left\{(U-c_1)^2\left[Y_z\right]^2-\left[(U-c_1)_z\right]^2Y^2\right\}\,dz \\
\delta &= \frac{1}{2c_0}\int_{-H}^{0}\rho_0Y^2\,dz \\
\mathcal{F} &= \int_{-H}^{0}\rho_0\left(U^2\right)_\eta Y_z\,dz + \rho_0(-H)[U(-H)]^2Y_z(-H)H_\eta
\end{aligned}$$

It should be noted that only γ and $\mathcal{F}$ are influenced by the basic flow. The expressions for other coefficients are fully equivalent to the expressions for the corresponding coefficients of the extended Korteweg-de Vries equation, used by Helfrich and Melville [4], where the basic flow were not considered. Note also that the forcing term of the amplitude equation derived by Grimshaw and Smyth [3] is just a particular case of our $\mathcal{F}$.

We conclude that for the evaluation of the coefficients of the equation (11) only the spectral problem (8) and the boundary problem (9) have to be solved. The first problem is a standard one and its numerical treatment is well developed. The second is not so standard because of its degeneracy, but it has a unique solution in the class of functions orthogonal to Y in $L^2[-H,0]$. This solution can be found numerically by the special choice of basis or by other known methods. As it is easily checked, any other solution produces the same value of the coefficients γ and $\mathcal{F}$.

References

[1] T. Taniuti, C.-C. Wei, *J. Phys. Soc. Japan*, **24**, 941-946 (1968).

[2] R. H. S. Grimshaw, *Stud. Appl. Math.*, **65**, 159-188 (1981).

[3] R. H. S. Grimshaw, N. Smyth, *J. Fluid Mech.*, **136**, 453-466 (1986).

[4] K. R. Helfrich, W. K Melville, *J. Fluid Mech.*, **167**, 285-308 (1984).